

UKCCNA Position Statement: Critical Care Nurse Staffing during Surge: September 2021

The UKCCNA acknowledges the sustained pressure critical care nurses (and other staff) continue to experience. We understand the significant negative impact this is having on nurses' physical and mental health and well-being.

Nurse-patient ratios

We set out our position on surge bedside nurse: patient ratios in our November 2020 and January 2021 position statements.^{1,2} During times of surge, the UKCCNA firmly believe patient safety and staff well-being is best secured if ratios are maintained at a minimum of one trained critical care nurse (TCCN)* and one registered healthcare professional** for two level 3 patients (compared with the normal 1:1 ratio^{3,4}), and one TCCN with one registered healthcare professional for every four level 2 patients (compared to the normal 1:2 ratio^{3,4}) as set out in National pandemic workforce guidance documents, such as that published by NHSE.⁵ Other support staff*** should also be available to assist with delivering patient care.

The UKCCNA position from 14th January 2021 remains unchanged. Further dilution of bedside nurse : patient ratios outside the guidance given in November 2020 should only happen in exceptional circumstances where there is a need to expand capacity despite escalation to regional and national critical care networks and when all local and regional mutual aid options, including inter-regional assistance, have been exhausted. Ratios beyond 1 TCCN : 2 patients should only be used to facilitate emergency admissions. If surge models are used to facilitate elective surgery, then extra nursing support must be in place. Ongoing work is exploring additions to the substantive establishment including Nursing Associates and Operating Department Practitioners, examining how they will interface into the current workforce model.

Ratios must be returned to normal as soon as practicable as these staffing levels are unsustainable for nurses' well-being and for patient safety. After episodes of surge, until further evidence is available, we must see a return to GPICS2 staffing model.³

The UKCCNA requests that the four Nations provide further guidance on how to manage the increased workforce requirements associated with additional expansion of critical care capacity. Data on adverse patient and staff outcomes should be collected.

Redeployment of staff

Surge ratios should be achieved through the redeployment of staff from outside critical care who should have received critical care surge training. Staff are encouraged to make use of the online training resources, such as those produced by the London Transformation and Learning

Collaborative - Critical Care (LTLC) - e-Learning for Healthcare (e-lfh.org.uk) and the RCN – Introducing Critical Care <https://www.rcn.org.uk/professional-development/professional-services/introducing-critical-care>.

Redeployment of staff, either within their own NHS organisation or between organisations is not always viable. This is due to multiple factors including staff sickness, pressures related to covering ward beds and the need for inter-hospital transfers. In addition, many staff redeployed during surge have been unable maintain their skills after return to their substantive posts and future redeployed staff may not be the same people.

This problem is compounded by the bureaucratic challenges faced by some nurses who do not currently work for the NHS (for example academics and retired staff) which prevent or delay them from being able to work. Unnecessary restrictions and administrative hurdles must be removed, and a common-sense approach taken to facilitate the timely availability of staff and the maintenance of fundamental critical care knowledge and skills, necessary for periods of surge, using resources such as the digital skills passport.

Paediatric patients

During an unseasonable Respiratory Syncytial Virus (RSV) surge, critically ill older paediatric patients may, on occasion, need to be cared for in adult ITU settings. Guidance around staffing requirements, ratios, education, and safeguarding can be found at: https://pccsociety.uk/wp-content/uploads/2016/05/PICS_standards_2015.pdf . Educational resources for Respiratory Surge in Children produced in collaboration with PCCS and HEE are also available on the [e-Learning for Healthcare](#) website.

Nurse well-being

The UKCCNA acknowledges the significant negative impact on nurses' physical and mental health and well-being. Professional Nurse Advocates should be given protected time to enable them to provide nurses with the much-needed support they require during and after experiencing unrelenting pressure to meet demands.

We strongly encourage nurses to take full advantage of available support resources such as the regional resilience hubs funded by NHSE. Asking for help is not a sign of weakness. Links to well-being resources are available via our member organisation websites: [British Association of Critical Care Nurses | BACCN](#); [Home | Intensive Care Society \(ics.ac.uk\)](#); [RCN - Home | Royal College of Nursing](#); [Coronavirus Resources & Guidance \(cc3n.org.uk\)](#). We also recommend that research exploring how best to support nurses' well-being is prioritised.

Definition of terms

***Trained critical care nurses (TCCN):** Whilst specialist training based on Steps competencies⁶ is usually required to become a TCCN, in this exceptional situation TCCNs are defined as those who have completed their step 1 competencies (or have been deemed competent to care for a critically ill patient) and who normally work in critical care plus TCCNs who have left the specialty within the last 12 months.

****Registered health care professionals:** A registered nurse (including nurse associates) or other relevant health care professionals (ideally with IV competence and some preparatory critical care training) working in the capacity of a bedside nurse. This will include nurses with critical care experience longer than 12 months ago.

*****Other support staff:** Healthcare support workers and non-nursing support workers, including students, dentists and doctors working in bedside care support roles.

The joint statement on developing immediate critical care nursing capacity remains current.⁷

<<<<<END>>>>>

Notes to readers:

The UK Critical Care Nursing Alliance (UKCCNA), established in 2013, provides a structured mechanism to facilitate collaborative working with all nationally recognised critical care nursing organisations across the United Kingdom. The aim of the UKCCNA is to be proactive and visionary about service requirements, providing quality assurance, enhancing the service, quality of care, patient experience and outcomes in critical care. Further details of UKCCNA member representatives are on the website: <https://www.ficm.ac.uk/UKCCNA>

The member organisations of the UKCCNA are:

- Royal College of Nursing (Critical Care and Flight Nursing Forum)
- British Association of Critical Care Nurses (BACCN)
- Critical Care National Network Nurse Lead Forum (CC3N)
- National Outreach Forum (NORF)
- Intensive Care Society (ICS)
- Paediatric Critical Care Society (PCCS)

References

1. UKCCNA position statement (Januray 2021): https://www.ficm.ac.uk/sites/default/files/ukccna_position_final_13.01.2021_.pdf
2. UKCCNA position statement (November 2020): <https://www.ficm.ac.uk/uk-critical-care-nursing-alliance-ukccna/ukccna-workforce-news-and-statements>
3. https://www.ics.ac.uk/ICS/ICS/GuidelinesAndStandards/GPICS_2nd_Edition.aspx
4. <https://www.england.nhs.uk/commissioning/spec-services/npc-crg/group-d/d05/>

5. <https://www.england.nhs.uk/coronavirus/publication/advice-on-acute-sector-workforce-models-during-covid-19/>
6. https://www.cc3n.org.uk/uploads/9/8/4/2/98425184/01_new_step_1_final_1_.pdf
7. <https://www.england.nhs.uk/coronavirus/publication/joint-statement-on-developing-immediate-critical-care-nursing-capacity/>

3rd September 2021.